

ORAL SESSION

ORAL SESSION 2B:

AGEING

Design and method: A sample of 521 subjects both genders, 253 centenarians (100.26 ± 1.98 years), and 268 controls (67.5 ± 13.25 years) subdivided into low (LCR, $n = 107$) and high (HCR, $n = 161$) CVR were included. Hypertension, diabetes, obesity (BMI, Kg.m^{-2}) and impaired kidney function, were defined according to standard criteria. CVR was calculated using Q risk 2–2016. DNA extraction was performed from the buccal epithelium sample using CitogeneâBuccal Kit, genotyping through iPLEX-MassARRAYâ, read by MALDI-TOF mass spectrometry and analyzed by EARTDECODEâ. Statistics were done throughout MDR and SPSS softwares. Significance defined as P-value < 0.05 .

Results: The variables that most were with the risk of not achieving exceptional longevity, OR (95% CI), were: BMI 1.558 (1.445–1.680), hypertension 2.358 (1.565–3.553), smoking-habits 4.528 (2.579–7.949), diabetes 5.553 (2.889–10.675), hypercholesterolemia 1.016 (1.010–1.022) and regular consumption of red meat 22.363 (13.987–35.755). There were differences in frequencies genotypic between centenarians and controls in ACE I/D, rs4646994 ($p = 0.001$) and NOS3 GT, rs1799983 ($p < 0.0001$) polymorphisms, being ACE DD and NOS3 GG genotypes more frequent in centenarians than in controls. No significant differences were observed concerning: AGT-rs699; AGT-rs4762; AGR1-rs5182; GRK4-rs2960306; GRK4-rs1024323; SLC12A3-rs13306673. There is a close genotype interaction between ACE and NOS3 each other and with hypertension, hypercholesterolemia, smoking and red meat intake. There is a predominance of Q risk marker (mainly environmental-related) associated with longevity in interaction to ACE I/D and NOS3.

Conclusions: In longevity pathway, even there is a gene-gene interaction particularly regarding ACE and NOS3, there is also increasing interaction between genes related to CVR, with the predominantly environmental factors that form the basis of Q risk determination. Underlie factors may modify gene expression modulated by the genotype and influence aging process.

GENETIC AND ENVIRONMENT INTERACTIONS CONTRIBUTE TO LONGEVITY: A CASE-CONTROL STUDY WITH CENTENARIANS

A. Pereira Da Silva Oliveira¹, L. Aguiar^{1,2}, A. Matos^{1,2}, J. Covas-Lima³, Â. Gil¹, J. Gorjão-Clara⁴, J. Polónia⁵, M. Bicho^{1,2,1} *Ecogenetics and Human Health Unit, Genetics Laboratory, Environmental Health Institut, ISAMB, Lisbon, PORTUGAL*, ²Instituto de Investigação Bento da Rocha Cabral, Lisbon, PORTUGAL, ³Coventry University, Faculty of Health and Life Sciences, Coventry, UNITED KINGDOM, ⁴University Geriatric Unit, Environmental Health Institute, ISAMB, Lisbon, PORTUGAL, ⁵Faculty of Medicine Oporto, Internal Medicine, Hypertension, Clinical Pharmacology, Porto, Portugal

Objective: Environmental or genetic factors, together with the interaction between them, may explain the increasing population of centenarians. This research aimed to understand these interactions that underlie centenarian's phenotypes, namely gene-gene and gene-environment, particularly in relation to cardiovascular risk (CVR).